






Uptake of an online psychological therapy program (iCanADAPT-Early) when implemented within a clinical pathway in cancer care centres

Fiona Davies¹  | Marnie Harris¹ | Joanne Shaw¹  | Phyllis Butow¹  |
 Jill Newby^{2,3} | Michael Murphy³ | Laura Kirsten^{1,4}  |
 The ADAPT Program Group¹ | Heather L. Shepherd^{1,5} 

¹School of Psychology, The University of Sydney, Psycho-Oncology Co-operative Research Group (PoCoG), Sydney, New South Wales, Australia

²School of Psychology, Faculty of Science, UNSW Sydney, Sydney, New South Wales, Australia

³Clinical Research Unit for Anxiety and Depression, UNSW Sydney, Sydney, New South Wales, Australia

⁴Nepean Cancer Care Centre, Sydney, New South Wales, Australia

⁵Sydney Nursing School, Faculty of Medicine and Health, The University of Sydney, Sydney, New South Wales, Australia

Correspondence

Heather L. Shepherd, Susan Wakil Health Building (D18), The University of Sydney, NSW 2006, Australia.

Email: heather.shepherd@sydney.edu.au

Funding information

Cancer Institute NSW, Grant/Award Number: 14/TPG/1-02

KEYWORDS

anxiety, cancer, depression, implementation, internet cognitive behavioural therapy, psycho-oncology

Key points

- Providing access to an online psychological therapy program in cancer care did not result in significant uptake and engagement by patients, or referrals by cancer care professionals
- Despite many opportunities for referral by cancer care professionals, only 2% resulted in a referral to the program. None of these referrals were taken up
- Twenty five patients self-referred, however only seven (28%) matched the recommended criteria for the program of mild to moderate psychological distress
- Patient engagement in the program was low, with 11/25 (44%) patients completing at least one lesson
- Cancer care professionals reported numerous barriers to use including patient barriers such as fatigue, and personal barriers such as forgetting to refer

1 | INTRODUCTION

Within cancer care, there has been increasing importance placed on the psychosocial well-being of patients. Meta-analysis indicates rates of depression of 5.4% to 49.0% (pooled prevalence: 21.0%) and for anxiety 3.4% to 43.0% (pooled prevalence: 21.0%) in

cancer patients, which is higher than in the general population.¹

Whilst some cancer services have established psycho-oncology services, workforce limitations, access issues, a focus on physical health as well as stigma associated with mental health, limit appropriate available support and uptake. Online cognitive behavioural therapy (CBT) for psychological symptoms can address

ADAPT Program group members: A/Prof Philip Beale, Prof Phyllis Butow, A/Prof Josephine Clayton, Jessica Cuddy, Dr Fiona Davies, A/Prof Haryana Dhillon, Dr Mona Faris, Liesbeth Geerligs, Prof Afaf Girgis, Dr Peter Grimison, Prof Tom Hack, Marnie Harris, Sharon He, Prof Brian Kelly, A/Prof Patrick Kelly, Dr Laura Kirsten, Dr Toni Lindsay, A/Prof Melanie Lovell, Dr Tim Luckett, Lindy Masya, Dr Michael Murphy, Dr Jill Newby, Don Piro, Dr Melanie Price, Dr Nicole Rankin, A/Prof Joanne Shaw, Prof Tim Shaw, Dr Heather Shepherd, Prof Rosalie Viney, Jackie Yim.

many of these barriers. However, a recent review of online therapies for cancer patients reported mixed results and concluded that while there is significant promise, there is currently inadequate evidence of their efficacy.² Evidence of post-trial implementation success, including patient uptake and engagement within usual cancer care settings, is even more limited.

An online transdiagnostic CBT program (iCanADAPT-Early) was developed to support implementation of a clinical pathway (ADAPT CP) for identification, assessment, referral and management of anxiety and depression in cancer patients.³ The ADAPT CP uses a stepped-care model, incorporating routine screening and referral and recommended treatment options. Patients complete brief screening tools via the web-based ADAPT Portal, to identify and quantify symptoms consistent with anxiety/depression. The clinical team are alerted and responsible for appropriate referral based on symptom severity. Online therapy is included within the ADAPT CP as a recommended option for patients with mild to moderate levels of anxiety and/or depression, to reduce the burden of care on the psycho-oncology service. The ADAPT CP implementation study was guided by the PARIHS implementation framework, which highlights the roles of context, facilitation and evidence in promoting sustainability.⁴

iCanADAPT-Early comprised an 8-lesson/16-week, clinician-monitored online cognitive-behavioural program, for patients with early stage cancer being treated or in survivorship, to treat clinically significant anxiety/depression. Based on a previously evaluated, effective online therapy for the general population, it was tailored for the cancer setting by an expert panel. Prior to each lesson, patients were screened for psychological distress with the K10,⁵ with those scoring above specified risk criteria followed up by the central clinician. Randomised controlled trial data demonstrated a significant decrease in symptoms in those receiving iCanADAPT-Early.⁶

iCanADAPT-Early was made available to 12 cancer services utilising the ADAPT CP as part of an implementation study.⁷ Staff could refer patients with mild to moderate anxiety/depression to iCanADAPT-Early, with patients also able to self-refer. All iCanADAPT-Early materials and details of the program structure were available to staff on the ADAPT Portal. Each psychosocial team was also contacted by the central iCanADAPT-Early clinician to provide information, answer any questions, and set up a contact protocol. Staff were informed of eligibility criteria (Step 3–4 anxiety/depression and stage 0–2/3 cancer) for iCanADAPT-Early but were not required to confirm staging. No reminders were provided, although these may have occurred informally. Here we report uptake of iCanADAPT-Early and staff reported barriers.

2 | METHODS

2.1 | Quantitative data

To describe referral, engagement, and iCanADAPT-Early uptake outcomes, ADAPT Portal screening events, patient ADAPT CP step

allocation and referrals made by staff to iCanADAPT-Early were collected, as well as information about who registered and enrolled in iCanADAPT-Early, K10 scores and lesson completion. These data enabled reporting on:

1. **Opportunities for referral** – number of screening events at Step 2 or 3 of the ADAPT CP.
2. **Referrals made** to iCanADAPT-Early – documented discussions with patients about iCanADAPT.
3. **Referral Uptake** – number of referred patients who enrolled in iCanADAPT-Early.
4. **Patient Engagement and Program Uptake** – number who completed at least one lesson of iCanADAPT-Early, and total number of lessons completed.
5. **Psychological distress** – ADAPT CP step level and K10 scores completed prior to each iCanADAPT-Early lesson.

2.2 | Qualitative

Staff perceptions of iCanADAPT-Early were captured during semi-structured audio-recorded interviews with staff familiar with the program.⁷ Interviews were transcribed verbatim and coded using NVivo12 Software. Transcripts were thematically analysed⁸ to identify barriers and facilitators to iCanADAPT-Early referral, uptake and engagement.

3 | RESULTS

3.1 | Study sample

Over the 12-month ADAPT CP implementation, there were 324 screening events which indicated Step 2 (mild) or Step 3 (moderate) levels of anxiety or depression; 188 at Step 2 and 136 at Step 3. Cancer staging reported showed 1 (0.3%), 37 (11%), 111 (34%) and 87 (27%) of screening events were by patients with a diagnosis of Stage 0, 1, 2/3, 4 cancer respectively. Staging was not reported for 88 screening events due to missing data or a haematological cancer diagnosis.

Thirteen staff participants consented and were interviewed. Ten (77%) were psychosocial staff, three (23%) were nursing staff. Participants had worked in cancer care an average of 5 years.

3.2 | Referrals to iCanADAPT

Only seven referrals (2%) from 324 screening events where iCanADAPT-Early was considered an appropriate referral option based on screening score and clinician recommendation were made. Clinician interviews highlighted barriers occurring at various stages including referral, uptake, and engagement. Barriers to referral included lack of program familiarity; forgetting to mention

iCanADAPT-Early as it was not part of routine care; perceived non-suitability of program for the patient due to their lack of technological expertise, illness or personality; a preference for more therapist involvement; or dislike of some format or content aspects of iCanADAPT-Early.

Twenty-five patients self-referred to iCanADAPT-Early, of whom seven (28%) were within the recommended anxiety/depression range based on their screening score (ADAPT CP Step 2 or 3, see Table 1). Thirteen (52%) screened at Step 1, which indicated no clinical need for additional psychological support.

3.3 | Referral uptake of iCanADAPT-Early

None of the seven referrals made within the ADAPT Portal to iCanADAPT-Early were taken up.

Staff participants reported perceived and actual barriers to referral uptake included patients prioritising the physical aspects of treatment, preference for face-to-face support, or patients feeling too unwell.

3.4 | Engagement with and uptake of the iCanADAPT-Early program lessons

Eleven (44%) of the 25 self-referred patients completed at least one lesson (Range 1–8, mean 3). Four (36%) screened within the recommended clinical range for iCanADAPT-Early. See Table 1. Initial K10 scores prior to lesson 1 reflected low distress too, with the majority ($n = 6$, 55%) showing a K10 score <20 (likely to be well), four (36%) scoring between 20 and 24 (mild distress), and one (9%) scoring over 30 (severe distress).

4 | DISCUSSION

Despite a successful efficacy study⁶ and initial enthusiasm for the program, iCanADAPT-Early was infrequently used across the 12 participating cancer services. We assessed success using several criteria: clinician referrals as a proportion of referral opportunities;

referral uptake by patients; appropriateness based on patients' levels of psychological distress; and patient engagement as indicated by lesson completion.

iCanADAPT-Early was not successful on any of these criteria. Referrals made by health professionals were low (less than 2.5% of referral opportunities), with no uptake of referrals by eligible patients. Most self-referred patients (52%) did not meet the recommended selection criteria, and most did not complete any lessons (56%).

Some identified barriers were specific to cancer care, but many aligned with a recent systematic review⁹ examining uptake of online therapies in general. Both in this review, and our study, barriers included patient factors (such as the patient being too unwell), health professional factors (such as forgetting to mention it), and organisational factors (minimal education of health professionals regarding the program), as well as concerns about online versus face-to-face treatment.

Low referral numbers by clinicians are striking, particularly given the implementation of routine screening during the ADAPT cluster RCT which facilitated identification of patients needing referral. Some staff may not have been comfortable with the model of care used in iCanADAPT-Early, where patients are monitored for safety concerns but do not receive any direct therapist support. It is unclear what the most acceptable model of care may be from the perspective of cancer care professionals, and we are currently conducting further research in this area to better understand their perspectives.

4.1 | Clinical implications

While the acceptability and effectiveness of iCanADAPT-Early was tested with cancer patients and iteratively revised, the implementation model was not the result of a co-design process involving staff or patients, or with consideration of alternative models of care. Our results reinforce the importance of co-design, as per guidelines available.^{9,10} This includes considering with stakeholders, the ideal format and content of the program, the optimal model of care to meld with existing clinical practice, educating clinicians on the program content, goals and evidence-base, marketing the program to

TABLE 1 ADAPT CP step allocation in the ADAPT Portal for patients who self-referred to iCanADAPT-Early and who completed ≥ 1 iCanADAPT-Early lesson

	Patient CP step allocation in the ADAPT portal						No step allocation	Total
	Step 1 ^a	Step 2	Step 3	Step 4	Step 5			
Number of patients who self-referred to iCanADAPT-Early (n)	13 (52%)	3 (12%)	4 (16%)	1 (4%)	0 (0%)	4 (16%)	25 (100%)	
Number of self-referred patients who completed ≥ 1 iCanADAPT-Early lesson (n)	6 (55%)	3 (27%)	1 (9%)	0 (0%)	0 (0%)	1 (9%)	11 (100%)	

^aSteps 1–5 are minimal, mild, moderate, severe, very severe anxiety/depression respectively.

patients, and training staff to make effective referrals which promote patient uptake.

5 | LIMITATIONS

The measure of clinician referrals relied on staff documenting referrals to iCanADAPT-Early in the ADAPT Portal. We do not know if this always occurred. However, feedback from interview participants was consistent with the low rate of referrals reported. Numbers offered iCanADAPT are too small to reliably assess uptake and a larger implementation study is required.

ACKNOWLEDGEMENTS

We acknowledge and thank the 12 cancer services, their clinical and administrative staff who participated in the cluster randomised trial. Participating services and staff included: Alan Coates Cancer Centre, Dubbo NSW, Tim Williams, Dr Florian Honeyball; Blacktown Cancer & Haematology Centre, Western Sydney LHD, Blacktown, NSW, Dr Purnima Sundaresan, Colette Cole; Chris O'Brien Lifehouse, Camperdown, NSW A/Prof Peter Grimison, Dr Toni Lindsay; Concord Cancer Centre, Concord Hospital, SLHD, A/Prof Philip Beale, Dr Alex Withers, Jessica Medd; Sydney Adventist Hospital, Dr Katharine Hodgkinson, Kate Wilson; Manning Hospital Cancer Care Services, Taree NSW, Rachel Pitt; Nepean Cancer Care Centre, Penrith NSW, Dr Laura Kirsten, Louise Maher, Betsy Sajish and Renee Grant; North West Cancer Centre, Tamworth Hospital, NSW A/Prof. Mathew George, Margaret Chamen, and Rebecca Griffiths; Prince of Wales Hospital, Head and Neck Cancer Service, Randwick, SESLHD, Dr Kerry Tiller, Penelope Burdekin, Dareneth Place; St George Cancer Centre, Kogarah, SESLHD, NSW, Natalie Katalinic, Alison Szwajcer, Moira Way, Pauline Thomson, Emma Hair, Bronwyn King; Sydney Cancer and Haematology Services, Royal Prince Alfred Hospital, SLHD John Chalmers, Hannah Blyth. The ADAPT Program is funded by a Translational Program Grant (14/TPG/1-02) from the Cancer Institute NSW. The funding bodies had no role in study design, data collection, analysis or writing of the manuscript.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

CLINICAL TRIAL REGISTRATION

The ADAPT Cluster RCT is registered with the ANZCTR Registration number: ACTRN12617000411347.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Fiona Davies  <https://orcid.org/0000-0002-5247-9231>

Joanne Shaw  <https://orcid.org/0000-0002-9543-7066>

Phyllis Butow  <https://orcid.org/0000-0003-3562-6954>

Laura Kirsten  <https://orcid.org/0000-0002-8520-3098>

Heather L. Shepherd  <https://orcid.org/0000-0002-3350-1325>

REFERENCES

1. Brandenburg D, Maass SWMC, Geerse OP, et al. A systematic review on the prevalence of symptoms of depression, anxiety and distress in long-term cancer survivors: Implications for primary care. *Eur J Canc Care*. 2019;28:e13086. doi:10.1111/ecc.13086
2. McAlpine H, Joubert L, Martin-Sanchez F, Merolli M, Drummond KJ. A systematic review of types and efficacy of online interventions for cancer patients. *Patient Educ Couns*. 2015;98:283-295. doi:10.1016/j.pec.2014.11.002
3. Butow P, Price MA, Shaw JM, et al. Clinical pathway for the screening, assessment and management of anxiety and depression in adult cancer patients: Australian guidelines. *Psycho-Oncology*. 2015; 24(9):987-1001. doi:10.1002/pon.3920
4. Kitson A, Harvey G, McCormack B. Enabling the implementation of evidence based practice: a conceptual framework. *Quality in Health care*. 1998;7(3):149-158.
5. Mroczke RKD. *The Kessler Psychological Distress Scale (K10)*. 1994.
6. Murphy M, Newby J, Butow P, et al. Randomised controlled trial of internet-delivered cognitive behaviour therapy for clinical depression and/or anxiety in cancer survivors (iCanADAPT Early). *Psycho-oncology*. 2020;29(1):76-85. doi:10.1002/pon.5267
7. Butow P, Shaw J, Shepherd HL, et al. Comparison of implementation strategies to influence adherence to the clinical pathway for screening, assessment and management of anxiety and depression in adult cancer patients (ADAPT CP): study protocol of a cluster randomised controlled trial. *BMC Canc*. 2018;18(1):1-12. doi:10.1186/s12885-018-4962-9
8. Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol*. 2008;8(1):1-10. doi:10.1186/1471-2288-8-45
9. Davies F, Shepherd HL, Beatty L, Clark B, Butow P, Shaw J. Implementing web-based therapy in routine mental health care: systematic review of health professionals' perspectives. *J Med Internet Res*. 2020;22(7):e17362. doi:10.2196/17362
10. Bevan Jones R, Stallard P, Agha SS, et al. Practitioner review: Co-design of digital mental health technologies with children and young people. *JCPP (J Child Psychol Psychiatry)*. 2020;61:928-940. doi:10.1111/jcpp.13258

How to cite this article: Davies F, Harris M, Shaw J, et al. Uptake of an online psychological therapy program (iCanADAPT-Early) when implemented within a clinical pathway in cancer care centres. *Psychooncology*. 2021;1-4. doi:10.1002/pon.5808